



FORTA-WEAN W/ CTC-DENA

MEDICATED

FOR STARTER PIGS 18 TO 25 POUNDS

For the control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin. Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENTS

Chlortetracycline	400 g/ton
Tiamulin HydrogenFumarate	35 g/ton

GUARANTEED ANALYSIS

Crude Protein	Min	21.00%		
L-Lysine	Min	1.40%		
Crude Fat	Min	4.50%		
Crude Fiber			Max	3.50%
Calcium(Ca)	Min	0.60	Max	1.10%
Phosphorus(P)	Min	0.65%		
Salt(NaCl)	Min	0.30	Max	0.90%
Selenium(Se)	Min	0.3 ppm		
Zinc(Zn)	Min	1950 ppm		

INGREDIENTS

Grain Products, Plant Protein Products, Lactose, Monocalcium Phosphate, Calcium Carbonate, Salt, Zinc Oxide, Yucca Schidigera Extract, Animal Protein Products, Vitamin E Supplement, Niacin Supplement, d-Calcium Pantothenate, Vitamin A Acetate, Riboflavin Supplement, Thiamine Mononitrate, Vitamin D₃ Supplement, Folic Acid, Dried *Bacillus Licheniformis* Fermentation Product, Dried *Bacillus subtilis* Fermentation Product, Menadione Sodium Bisulfite Complex, Biotin, Vitamin B₁₂ Supplement, Pyridoxine Hydrochloride, L-Tryptophan, L-Threonine, Copper Sulfate, Magnesium Oxide, Choline Chloride, Ferrous Sulfate, Manganous Oxide, Sodium Selenite, Ethylenediamine Dihydriodide, Benzoic Acid, Citric Acid, Sorbic Acid, Sodium Benzoate, BHA & BHT(as preservatives), Formic Acid, Lactic Acid, Fumaric Acid, Sodium Hydroxide, L-Lysine, Potassium Chloride, DL-Methionine, Natural & Artificial Flavors

FEEDING DIRECTIONS

Feed as a sole ration to pigs weighing 18 to 25 lbs. body weight. Feed continuously as sole ration for 14 days.

WARNING: Withdraw 2 days before slaughter.

Caution: Do not feed undiluted. Do not feed to animals other than swine. The effects of Tiamulin on swine reproductive performance, pregnancy and lactation have not been determined. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

MANUFACTURED BY:
LOWE'S PELLETS & GRAIN, INC.
GREENSBURG, IN 47240
NET WEIGHT 50 LBS. (22.68 KG) OR BULK

LAST UPDATED
12302013 Green

6.29.2018