



# BOAR RATION

## MEAL W/ PULMOTIL

### #20810

#### BOAR MAINTENANCE RATION

For increased rate of weight gain and improved efficiency, grower feeds.

#### ACTIVE DRUG INGREDIENT

TILMICOSIN

181 G/TON

#### GUARANTEED ANALYSIS

Crude Protein	Min	16.00%		
L-Lysine	Min	0.90%		
Crude Fat	Min	3.40%		
Crude Fiber			Max	5.00%
Calcium(Ca)	Min	0.65%	Max	1.15%
Phosphorus(P)	Min	0.50%		
Salt (NaCl)	Min	0.20%	Max	0.70%
Selenium(Se)	Min	0.3 ppm		
Zinc(Zn)	Min	180 ppm		
Phytase		435 FTU/lb.		

#### INGREDIENTS

Grain Products, Plant Protein Products, Roughage Products, Monocalcium Phosphate, Calcium Carbonate, Salt, Animal & Vegetable Fat (preserved with Ethoxyquin), L-Lysine, Dried Schizosaccharomyces pombe Fermentation Product, Potassium Sulfate, Magnesium Sulfate, Sodium Sulfate, Psyllium Seed Husks, Vitamin A Acetate in Gelatin, Vitamin D3 Supplement, Vitamin E Supplement, Vitamin B12 Supplement, Pyridoxine Hydrochloride, Thiamine Mononitrate, Menadione Sodium Bisulfite Complex, Riboflavin Supplement, d-Calcium Pantothenate, Niacin Supplement, Choline Chloride, Folic Acid, Zinc Oxide, Ferrous Sulfate, Zinc Sulfate, Manganous Oxide, Biotin, Ascorbic Acid, Copper Sulfate, Ethylenediamine Dihydriodide, Selenium Yeast, Sodium Selenite, Zinc Amino Acid Complex, Iron Amino Acid Complex, Copper Amino Acid Complex.

#### FEEDING DIRECTIONS

Feed 4 to 6 pounds per day to maintain desired condition.

#### WARNING

Withdraw 7 days prior to slaughter.

#### LIMITATIONS FOR USE

Federal law limits this drug to use under the professional supervision of a licensed veterinarian. Animal feed bearing or containing this veterinarian feed directive drug shall be fed to animals only by or upon a lawful veterinary feed directive issued by a licensed veterinarian in the course of the veterinarian's professional practice. Feed continuously as the sole ration for a 21 day period, beginning approximately 7 days before an expected disease outbreak. Feed containing Tilmicosin shall not be feed to pigs for more than 21 days during each phase of production without ceasing administration for re-evaluation of antimicrobial use by a licensed veterinarian before reinitiating further course of therapy with an appropriate antimicrobial.

#### CAUTION

Do not allow horses or other equines access to feeds containing Tilmicosin. The safety of Tilmicosin has not been established in male swine intended for breeding purposes. Do not use in feeds containing bentonite. Bentonite in feeds may affect the efficacy of Tilmicosin.

MANUFACTURED BY  
LOWE'S PELLETS & GRAIN, INC.  
GREENSBURG, IN 47240  
NET WEIGHT 50 LBS. (22.68 KG) OR BULK

6.29.2018

LOT# |