



**15% GOAT W/RUM PELLETT  
MEDICATED  
#3015**

**NEW SMALLER PELLETT  
FOR IMPROVED PERFORMANCE**

**Complete pelleted feed for growing & finishing goats.**

For the prevention of coccidiosis caused by *Eimeria crandallis*, *Eimeria christenseni* and *Eimeria ninakohlyakimovae*.

**ACTIVE DRUG**

MONENSIN ..... 20G/TON

**GUARANTEED ANALYSIS**

CRUDE PROTEIN	MIN	15.0%		
(This product contains not more than .8% equivalent crude protein from non-protein nitrogen)				
CRUDE FAT	MIN	3.50%		
CRUDE FIBER			MAX	14.0%
ACID DETERGENT FIBER			MAX	16.0%
CALCIUM	MIN	0.50	MAX	1.00%
PHOSPHORUS	MIN	0.40%		
SALT	MIN	0.50	MAX	1.00%
COPPER (CU)	MIN	30 PPM	MAX	40 PPM
SELENIUM	MIN	0.70 PPM		
VITAMIN A	MIN	6250 IU/lb		

**INGREDIENTS**

GRAIN PRODUCTS, PROCESSED GRAIN BY-PRODUCTS, ROUGHAGE PRODUCTS, ALFALA PRODUCTS, PORCINE BLOOD MEAL, FISH MEAL, CALCIUM CARBONATE, MOLASSES PRODUCTS, SALT, AMMONIUM CHLORIDE, YEAST CULTURE (SACCHAROMYCES CEREVISIAE YEAST), VITAMIN E SUPPLEMENT, MAGNESIUM SULFATE, POTASSIUM SULFATE, POTASSIUM CHLORIDE, MAGNESIUM OXIDE, VITAMIN A SUPPLEMENT, VITAMIN D3 SUPPLEMENT, COPPER SULFATE, COBALT CARBONATE, MANGANESE SULFATE, ZINC SULFATE, ETHYLENEDIAMINE DIHYDRIDIDE, FERROUS SULFATE, SODIUM SELENITE.

**CAUTION: Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Consumption by unapproved species may result in toxic reaction. Do not feed to lactating goats.**

**WARNING: DO NOT FEED TO SHEEP. THIS PRODUCT CONTAINS COPPER.**

**FEEDING DIRECTIONS:** Feed continuously as the sole concentrate to growing and finishing goats with 40% to 70% of the diet as a good quality forage.

MANUFACTURED BY  
LOWE'S PELLETS & GRAIN, INC.  
GREENSBURG , IN 47240  
NET WEIGHT 50 LBS. (22.68 KG) OR BULK

PINK 4.25.2019 LOT[63429 |