



## Beef Commander w/ Rumensin

### #4550

### MEDICATED

A Protein Supplement for Beef Cattle.

For improved feed efficiency.

#### ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) ..... 250.0 g/ton

#### GUARANTEED ANALYSIS

Crude Protein	(Min)		35.0%
(This includes not more than 15.1% equivalent non-protein nitrogen)			
Crude Fat	(Min)		3.0 %
Crude Fiber	(Max)		6.5 %
Calcium	(Min)	5.0	(Max) 5.5 %
Phosphorus	(Min)		0.6 %
Salt	(Min)	3.6	(Max) 4.6 %
Potassium	(Min)		1.0 %
Zinc	(Min)		900.0 ppm
Copper	(Min)		280.0 ppm
Selenium	(Min)		2.9 ppm
Vitamin A	(Min)		50,000.0 IU/lb
Vitamin E	(Min)		50.0 IU/lb

#### LIST OF INGREDIENTS

Plant Protein Products, Processed Grain By-Products, Calcium Carbonate, Calcium Chloride, Urea, Roughage Products, Salt, Magnesium Oxide, Animal Fat, Sodium Bentonite, Potassium Chloride, Vitamin A Supplement, Copper Sulphate, Sodium Selenite, Vitamin D3 Supplement, Vitamin E Supplement, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydriodide, Calcium Iodate, Cobalt Sulfate, Mineral Oil, Cobalt Carbonate.

#### FEEDING DIRECTIONS

Feed at a rate of 2 lbs./head/day to provide 250 mg/head/day of Monensin (Rumensin®). Provide abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further directions consult your supplier or nutritionist.

**WARNING:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. **CAUTION:** Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. **CAUTION:** This feed contains added copper, do not feed to sheep. **CAUTION:** Do not exceed 0.3 mg/kg of selenium in the complete feed.

MANUFACTURED BY  
LOWE'S PELLETS & GRAIN, INC.  
Greensburg, IN 47240  
**(800) 888-7863**  
NET WT 50 LB (22.68 kg) or BULK

YELLOW

4.26.2019

LOT#63384