



**EXCALIBUR II**  
**PELLET W/RUM MEDICATED**  
**#3006**

**Complete pelleted feed to be used as an initial creep feed and starter for goats from birth to six months, as well as a developer for maximizing the genetic potential of the show buck and doe.**

For the prevention of coccidiosis caused by Eimeria crandallis, Eimeria christenseni and Eimeria ninakohlyakimovae.

**ACTIVE DRUG**

MONENSIN .....20G/TON

**GUARANTEED ANALYSIS**

CRUDE PROTEIN	MIN	15.0%		
(This product contains not more than .8% equivalent crude protein from non-protein nitrogen)				
CRUDE FAT	MIN	4.25%		
CRUDE FIBER			MAX	16.0%
ACID DETERGENT FIBER			MAX	18.0%
CALCIUM	MIN	1.10%	MAX	1.60%
PHOSPHORUS	MIN	0.35%		
SALT	MIN	0.80%	MAX	1.30%
COPPER (CU)	MIN	10 PPM	MAX	30 PPM
SELENIUM	MIN	0.300 PPM		
VITAMIN A	MIN	16,000 IU/lb		
VITAMIN D	MIN	2,500 IU/lb.		
VITAMIN E	MIN	50 IU/lb		

**INGREDIENTS**

ROLLED CORN, SOYBEAN HULLS, SUNCURED ALFALFA MEAL, SOYBEAN MEAL, WHEAT MIDDS, CANOLA MEAL, STABILIZED RICE BRAN, LINSEED MEAL, MOLASSES PRODUCTS, CALCIUM CARBONATE, ANIMAL & VEGETABLE FAT, SALT, YEAST CULTURE (SACCHAROMYCES CEREVISIAE YEAST), AMMONIUM CHLORIDE, THIAMINE, WHEAT BRAN, DRIED ASPERGILLUS ORYZAE FERMENTATION PRODUCTS, ZINC METHIONINE COMPLEX, ZINC AMINO ACID COMPLEX, MANGANESE AMINO ACID COMPLEX, COPPER AMINO ACID COMPLEX, COBALT GLUCOHEPTONATE, VITAMIN E SUPPLEMENT, MAGNESIUM OXIDE, VITAMIN A SUPPLEMENT, VITAMIN D3 SUPPLEMENT, COPPER SULFATE, COBALT CARBONATE, MANGANESE SULFATE, ZINC SULFATE, ETHYLENEDIAMINE DIHYDRIDIDE, FERROUS SULFATE, SODIUM SELENITE.

**FEEDING DIRECTIONS:**

- Feed Excalibur Goat continuously at a rate of 3% to 4% of body weight to all classes of goats for added bloom and to stimulate growth rate.
- Feed Excalibur Goat continuously at a rate of 1% to 2% of body weight to mature bucks and does to maintain bloom and freshness of muscle expression.

**CAUTION:** Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Consumption by unapproved species may result in toxic reaction. Do not feed to lactating goats.

**WARNING: DO NOT FEED TO SHEEP THIS PRODUCT CONTAINS COPPER.**

MANUFACTURED BY  
LOWE'S PELLETS & GRAIN, INC.  
GREENSBURG , IN 47240  
NET WEIGHT 50 LBS. (22.68 KG) OR BULK



ORANGE

9.18.2019

LOT# 66828